

Biostatistics 140.623 Third Term, 2002-2003

Laboratory Exercise 5

Below find times to “drug failure” (as determined by a treating psychiatrist) for 15 patients in a study comparing a new treatment for schizophrenia to a standard treatment (modification of SEP #11).

Trt group	Times (wks)
Standard	3, 5+, 6+, 9, 13+, 15+, 16+
New	4, 6, 9, 9, 10+, 11+, 13+, 14+

1. Construct the Kaplan-Meier survival curves by treatment. Compare to the Stata log on the next page:

Standard Treatment					New Treatment				
Event-Time (t_i)	Number at Risk (n_i)	Events (y_i)	$\frac{(n_i - y_i)}{n_i}$	$\hat{S}(t_i)$	Event-Time (t_i)	Number at Risk (n_i)	Events (y_i)	$\frac{(n_i - y_i)}{n_i}$	$\hat{S}(t_i)$

. list

	weeks	trt	id	failure
1.	3	0	1	1
2.	5	0	2	0
3.	6	0	3	0
4.	9	0	4	1
5.	13	0	5	0
6.	15	0	6	0
7.	16	0	7	0
8.	4	1	8	1
9.	6	1	9	1
10.	9	1	10	1
11.	9	1	11	1
12.	10	1	12	0
13.	11	1	13	0
14.	13	1	14	0
15.	14	1	15	0

```
. stset weeks, failure(failure==1) id(id)
```

```

          id: id
      failure event: failure == 1
obs. time interval: (weeks[_n-1], weeks]
exit on or before: failure

```

```
-----
15 total obs.
0 exclusions
-----
```

```

15 obs. remaining, representing
15 subjects
6 failures in single failure-per-subject data
143 total analysis time at risk, at risk from t = 0
      earliest observed entry t = 0
      last observed exit t = 16

```

```
. sts list if trt==0
```

```

      failure _d: failure == 1
analysis time _t: weeks
          id: id

```

Time	Beg. Total	Fail	Net Lost	Survivor Function	Std. Error	[95% Conf. Int.]	
3	7	1	0	0.8571	0.1323	0.3341	0.9786
5	6	0	1	0.8571	0.1323	0.3341	0.9786
6	5	0	1	0.8571	0.1323	0.3341	0.9786
9	4	1	0	0.6429	0.2104	0.1515	0.9017
13	3	0	1	0.6429	0.2104	0.1515	0.9017
15	2	0	1	0.6429	0.2104	0.1515	0.9017
16	1	0	1	0.6429	0.2104	0.1515	0.9017


```
. sts list if trt==1
```

```

failure_d: failure == 1
analysis time t: weeks
id: id
    
```

Time	Beg. Total	Fail	Net Lost	Survivor Function	Std. Error	[95% Conf. Int.]	
4	8	1	0	0.8750	0.1169	0.3870	0.9814
6	7	1	0	0.7500	0.1531	0.3148	0.9309
9	6	2	0	0.5000	0.1768	0.1520	0.7749
10	4	0	1	0.5000	0.1768	0.1520	0.7749
11	3	0	1	0.5000	0.1768	0.1520	0.7749
13	2	0	1	0.5000	0.1768	0.1520	0.7749
14	1	0	1	0.5000	0.1768	0.1520	0.7749

2. Based upon the plot of the Kaplan-Meier curves for each treatment group, which treatment, if any, should be preferred?



3. Calculate the log-rank statistic to test whether overall drug failure differs between the two treatments. Compute by hand the log-rank test statistic from the 2x2 tables based on each event time.

$$\chi^2_{LR} = \frac{\left[\sum_j (a_j - E(a_j)) \right]^2}{\sum_j \hat{V}ar(a_j)} \text{ where } E(a_j) = \frac{d_j n_{ja}}{n_j} \text{ and } \hat{V}ar(a_j) = \frac{d_j (n_j - d_j) n_{ja} n_{jb}}{n_j^2 (n_j - 1)}$$

	Event	No Event	Total
Standard Trt	a_j		n_{ja}
New Trt	c_j		n_{jb}
Total	d_j		n_j

	Event	No Event	Total
Standard Trt			
New Trt			
Total			

	Event	No Event	Total
Standard Trt			
New Trt			
Total			

	Event	No Event	Total
Standard Trt			
New Trt			
Total			

	Event	No Event	Total
Standard Trt			
New Trt			
Total			

Compare your calculation to that obtained by Stata below.

Log-rank test for equality of survivor functions

trt	Events observed	Events expected
0	2	2.51
1	4	3.49
Total	6	6.00

chi2(1) = 0.20
 Pr>chi2 = 0.6531